

Fachspezifische lesenswerte Literatur

Pauley T, Devlin M, Madan-Sharma P. A single-blind, cross-over trial of hip abductor strength training to improve timed up & go performance in patients with unilateral, transfemoral amputation. *J Rehabil Med* 2014;46:XX-XX

Tinl ML, Kale MK, Doshi S, Guarino AJ, Beninato M. The mobility scale for acute stroke (MSAS) predicts discharge destination after acute hospitalization. *J Rehabil Med* 2014;46:00-00

Limbäck G, Wendt GK, Thomeé. A structured physiotherapy treatment model can provide rapid relief to patients who qualify for lumbar disc surgery: a prospective cohort study. *J Rehabil Med* 2014;46:00-00

Fonseca H, Moreira-Goncalves D, Coriolano HJA, Duarte JA. Bone quality: the determinants of bone strength and fragility. *Sports Med* 2013; DOI 10.1007/s40279-013-0100-7

Murphy SL, Alexander NB, Levoska M, Smith DM. Relationship between fatigue subsequent physical activity among older adults with symptomatic osteoarthritis. *Arthrit Care Res* 2013;65:1617-24

Grudyte-Racine R, Erlandson MC, Jackowski SA, Baxter-Jones ADG. Structural strength development at the proximal femur in 4-to 10-year-old precompetitive gymnasts: a 4-year longitudinal hip structural analysis study. *J Bone Miner Res* 2013;28:2592-600

Morgan P, McGinley J. Gait function and decline in adults with cerebral palsy: a systematic review. *Disabil Rehabil* 2014;36:1-9

Ploumis A, Varvarousis D, Konitsiotis S, Beris A. Effectiveness of botulinum toxin injection with and without needle electromyographic guidance for the treatment of spasticity in hemiplegic patients: a randomized controlled trial. *Disabil Rehabil* 2014;36:313-8

Cantarero-Villanueva I, Fernández-Lao C, Cuesta-Vargas AI, Del Moral-Avila R, Fernández-de-las-Penas C, Arroyo-Morales M. The effectiveness of deep water aquatic exercise program in cancer-related fatigue in breast cancer survivors: a randomized controlled trial. *Arch Phys Med Rehabil* 2013; <http://dx.doi.org/10.1016/j.apmr.2012.09.008>

Ma C, Szeto GP, Yan T, Wu S, Lin C, Li L. Comparing biofeedback with active exercise and passive treatment for the management of work-related neck and shoulder pain: a randomized controlled trial. *Arch Phys Med Rehabil* 2011;92:849-58

Hagen S, Stark D, Glazener C, Dickson S, Barry S, Elders A, Frawley H, Galea MP, Logan J, McDonald A, McPherson G, Moore KH, Norrie J, Walker A, Wilson D, on behalf of the POPPY Trial Collaborators. Individualised pelvic floor muscle training in women with pelvic organ prolapse (POPPY): a multicentre randomised controlled trial. *Lancet* 2014;383:796-806

Latham NK, Harris BA, Bean JF, Heeren T, Goodyear C, Zawacki S, Heislein DM, Mustafa J, Pardasaney P, Giorgetti M, Holt N, Goehring L, Jette AM. Effect of a home-based program on functional recovery following rehabilitation after hip fracture: a randomized clinical trial. *JAMA* 2014;311:700-8

Dardas A, Bae GH, Yule A, Wright J, Straughn N, Day CS. Acetic acid iontophoresis for recalcitrant scarring in post-operative hand patient. *J Hand Ther* 2014;27:44-8

Smoot B, Boyd BS, Byl N, Dodd M. Mechanosensitivity in the upper extremity following breast cancer treatment. *J Hand Ther* 2014;27:4-11

Oiestad BE, Holm I, Aune AK, Gunderson R, Myklebust G, Engebretsen L, Fosdahl MA, Risberg MA. Knee function and prevalence of knee osteoarthritis after anterior cruciate ligament reconstruction: a prospective study with 10 to 15 years of follow-up. *Am J Sports Med* 2010;38:2201

Mataran-Penarrocha GA, Castro-Sanchez AM, Garcia GC, Moreno-Lorenzo C, Carreno TP, Zafra MDO. Influence of craniosacral therapy on anxiety, depression and quality of life in patients with fibromyalgia. Evidence-Based Complementary and Alternative Medicine 2011; Article ID 178769, DOI: 10.1093/ecam/nep125

Reid SA, Rivett DA, Katekar MG, Callister R. Sustained natural apophyseal glides (SNAGs) are an effective treatment for cervicogenic dizziness. *Manual Ther* 13:357-66

Stone EE, Skubic M. Passive in-home measurement of stride-to-stride gait variability compared with vision and Kinect sensing. 23rd Annual International Conference of the IEEE EMBS Boston, Massachusetts USA, August 30 – September 3, 2011. 978-1-4244-4122-8/11 IEEE EMBS.

Wang F, Skubic M, Rantz M, Cuddihy PE. Quantitative gait measurement with pulse-doppler radar for passive in-home gait assessment. DOI 10.1109/TBME.2014.2319333, *IEEE Transactions on Biomedical Engineering*.

Lee HK, Cho SP, Xou JH, Lee KJ. The concurrent validity of the body center of mass in accelerometric measurement. Proceeding of the 20th Annual International Conference of the IEEE EMBS, Cité Internationale, Lyon, France August 23-26, 2007, 1-4244-0788-5/07 IEEE.

Clarsen B, Bahr R, Andersson SH, Munk R, Myklebust G. Reduced glenohumeral rotation, external rotation weakness and scapular dyskinesis are risk factors for shoulder injuries among elite male handball players: a prospective cohort study. *Br J Sports Med* 2014;48:1327-33

Otoshi K, Takegami M, Sekiguchi M, Onishi Y, Yamazaki S, Otani K, Shishido H, Kikuchi S, Konno S. Association between kyphosis and subacromial impingement syndrome: LOHAS study. *J Shoulder Elbow Surg* 2014; <http://dx.doi.org/10.1016/j.jse.2014.04.010>

Leung KS, Li CY, Tse YK, Choy TK, Leung PC, Hung VWY, Chan SY, Leung AHC, Cheung WH. Effects of 18-month low-magnitude high-frequency vibration on fall rate and fracture risks in 710 community elderly—a cluster-randomized controlled trial. *Osteoporos Int* 2014;25:1785-95

Haakstad LAH, Bo K. Effect of a regular exercise programme on pelvic girdle and low back pain in previously inactive pregnant women: a randomized controlled trial. *J Rehabil Med* 2015;47:00-00, DOI: 10.2340/16501977-1906

Naylor JM, Crosbie J, Ko V. Is there a role for rehabilitation streaming following total knee arthroplasty? Preliminary insights from a randomized controlled trial. *J Rehabil Med* 2015;47:00-00, DOI: 10.2340/16501977-1919

Aboagye E, Karlsson ML, Hagberg J, Jensen I. Cost-effectiveness of early interventions for non-specific low back pain: a randomized controlled study investigating medical yoga, exercise therapy and self-care advice. *J Rehabil Med* 2015;47:00-00, DOI: 10.2340/16501977-1910

Ng SSM, Cheung SY, Lai LSW, Liu ASL, Leong SHI, Fong SSM. Five times sit-to-stand test completion times among older women: influence of seat height and arm position. *J Rehabil Med* 2015;47:00-00, DOI: 10.2340/16501977-1915

Marusiak J, Zeligowska E, Mencil J, Kisiel-Sajewicz K, Majerczak J, Zoladz JA, Jaskolski A, Jaskolska A. Interval training-induced alleviation of rigidity and hypertonia in patients with Parkinson's disease is accompanied by increased basal serum brain-derived neurotrophic factor. *J Rehabil Med* 2015;47:00-00, DOI: 10.2340/16501977-1931

Marcocci C, Cetani F. Primary hyperparathyroidism. *N Engl J Med* 2011;365:2389-97

Terkelsen AJ, Molgaard H, Hansen J, Finnerup NB, Kroner K, Jensen TS. Heart rate variability in complex regional pain syndrome during rest and mental and orthostatic stress. *Anesthesiology* 2012;116:133-46

Morone G, Paolucci T, Alcuri MR, Vulpiani MC, Matano A, Bureca I, Paolucci S, Saraceni VM. Quality of life improved by multidisciplinary back school program in patients with chronic non-specific low back pain: a single blind randomized controlled trial. *Eur J Phys Rehabil Med* 2011;47:533-41

Tüzün S, Aktas I, Akarirmak Ü, Sipahi S, Tüzün F. Yoga might be an alternative training for the quality of life and balance in postmenopausal osteoporosis. *Eur J Phys Rehabil Med* 2010;46:69-72

Larossa F, Dura MJ, Menacho J, Gonzalez-Sabate L, Cordon A, Hernandez A, Garcia-Ibanez L. Aphysiologic performance on dynamic posturography in work-related patients. *Eur Arch Otorhinolaryngol* 2012; DOI 10.1007/s00405-012-1930-x

Custodio CM. Electrodiagnosis in cancer treatment and rehabilitation. *Am J Phys Med Rehabil* 2011;90:S38-S49

Kim SH, Kim M, Ahn YB, Lim HK, Kang SG, Cho JH, Park SJ, Song SW. Effect of dance exercise on cognitive function in elderly patients with metabolic syndrome: a pilot study. *J Sports Sci Med* 2011;10:671-8

Souza JC, Simoes HG, Campell CSG, Pontes FL, Boullosa DA, Prestes J. Haemophilia and Exercise. *Int J Sports Med* 2012;33:83-8

Honaker JS, Forston MR, Davis EA, Wiesner MM, Morgan JA. Effects of non contact low-frequency ultrasound on healing of suspected deep tissue injury: a retrospective analysis. *Int Wound J* 2012;DOI: 10.1111/j.1742-481X.2012.00944.x

Williams S, Whatman C, Hume PA, Sheerin K. Kinesio taping in treatment and prevention of sports injuries. *Sports Med* 2012;42:153-64

Tsai HJ, Hung HC, Yang JL, Huang CS, Tsauo JY. Could kinesio tape replace the bandage in decongestive lymphatic therapy for breast-cancer-related lymphedema? A pilot study. *Support Care Cancer* 2009;17:1353-60

Hughes K, Bellis MA, Jones L, Wood S, Bates G, Eckley L, NcCoy E, Mikton C, Shakespeare T, Officer A. Prevalence and risk of violence against adults with disabilities: a systematic review and meta-analysis of observational studies. *Lancet* 2012;379:1621-29

Carr AJ, Robertsson O, Graves S, Price AJ, Arden NK, Judge A, Beard DJ. Knee replacement. *Lancet* 2012;379:1331-40

Balagué F, Mannion AF, Pellisé F, Cedraschi C. Non-specific low back pain. *Lancet* 2012;379:482-91

White DK, Keysor JJ, Neogi T, Felson DT, LaValley M, Gross KD, Niu J, Nevitt M, Lewis CE, Torner J, Fredman L. When it hurts, a positive attitude may help: association of positive affect with daily walking in knee osteoarthritis. Results from a multicenter longitudinal cohort study. *Arthritis Care Res* 2012;64:1312-9

Chung KC, Burns PB, Kim HM, Burke FD, Wilgis EFS, Fox DA. Long-term followup for rheumatoid arthritis patients in a multicenter outcomes study of silicone metacarpophalangeal joint arthroplasty. *Arthritis Care Res* 2012;64:1292-300

Macfarlane GJ, Paudyal P, Doherty M, Ernst E, Lewith G, MacPherson H, Sim J, Jones GT, Arthritis Research UK Working Group on Complementary and Alternative Therapies for the Management of the Rheumatic Diseases. A systematic review of evidence for the effectiveness of practitioner-based complementary and alternative therapies in the management of rheumatic diseases: rheumatoid arthritis. *Rheumatology* 2012;51:1707-13

Rizzoli R, Adachi JD, Cooper C, Dere W, Devogelaer JP, Diez-Perez A, Kanis JA, Laslop A, Mitlak B, Papapoulos S, Ralston S, Reiter S, Werhya G, Reginster JY. Management of glucocorticoid-induced osteoporosis. *Calcif Tissue Int* 2012;91:225-43

10. KREMS KONFERENZ, Management der Spastizität – Update therapeutischer Strategien, 13.11.2012, Donau-Universität Krems, Department für Klinische Neurowissenschaften und Präventionsmedizin, Programm & Abstracts, www.donau-uni.ac.at/neuro

Guirand T, Granger R, Gremeaux V, Bousquet M, Richard L, Soukarié L, Babin T, Labrunée M, Sanguignol F, Bosquet L, Golay A, Pathak A. Telephone support oriented by accelerometric measurements enhances adherence to physical activity recommendations in noncompliant patients after a cardiac rehabilitation program. *Arch Phys Med Rehabil* 2012;93:2141-7

DiSotto-Monastero M, Chen X, Fisch S, Donaghy S, Gomez M. Efficacy of 7 days per week inpatient admissions and rehabilitation therapy. *Arch Phys Med Rehabil* 2012;93:2165-9

Lamberg EM, Hagins M. The effects of low back pain on natural breath control during a lowering task. *Eur J Appl Physiol* 2012;112:3519-24

Kim Y, Son J, Yoon B. Intensive unilateral neuromuscular training on non-dominant side of low back improves balanced muscle response and spinal stability. *Eur J Appl Physiol* 2012; DOI 10.1007/s00421-012-2513-7

Terblanche E, Millen AME. The magnitude and duration of post-exercise hypotension after land and water exercises. *Eur J Appl Physiol* 2012; DOI 10.1007/s00421-012-2398-5

Lange M, Krohn-Grimberghe B, Petermann F. Mittelfristige Effekte einer multimodalen Behandlung bei Patienten mit Fibromyalgiesyndrom. *Schmerz* 2011; DOI 10.1007/s00482-010-1003-2

Salomonowitz G, Friedrich M, Güntert BJ. Medizinische Relevanz von Magnetfeldern zur Schmerztherapie. *Schmerz* 2011; DOI 10.1007/s00482-010-1005-0

Villareal DT, Chode S, Parimi N, Sinacore DR, Hilton T, Armamento-Villareal R, Napoli N, Qualls C, Shah K. Weight loss, exercise, of both and physical function in obese older adults. *N Engl J Med* 2011;364:1218-29

Herridge MS, Tansey CM, Matté A, Tomlinson G, Diaz-Granados N, Cooper A, Guest CB, Mazer D, Slutsky AS, Cheung AM. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med* 2011;364:1293-304

Watson JM, Kang'ombe AR, Soares MO, Chuang LH, Worthy G, Bland JM, Iglesias C, Cullum N, Torgerson D, Nelson EA. Use of weekly, low dose, high frequency ultrasound for hard to heal venous leg ulcers: the VenUS III randomised controlled trial. *BMJ* 2011;342:d1092; DOI 10.1136/bmj.d1092

Gary RA, Cress E, Higgins MK, Smith AL, Dunbar SB. Combined aerobic and resistance exercise program improves task performance in patients with heart failure. *Arch Phys Med Rehabil* 2011;92:1371-81

Cunningham JE, Kelechi T, Sterba K, Barthelemy N, Falkowski P, Chin SH. Case report of a patient with chemotherapy-induced peripheral neuropathy treated with manual therapy (massage). *Support Care Cancer* 2011;19:1473-6

Laird BJA, Walley J, Murray GD, Clausen E, Colvin LA, Fallon MT. Characterization of cancer-induced bone pain: an exploratory study. *Support Care Cancer* 2011;19:1393-401

Krohn M, Listing M, Tjahjono G, Reissbauer A, Peters E, Klapp BF, Rauchfuss M. Depression, mood, stress, and Th1/Th2 immune balance in primary breast cancer patients undergoing classical massage therapy. *Support Care Cancer* 2011;19:1303-11

Rayegani SM, Bahrami MH, Samadi B, Sedighipour L, Mokhtarirad MR, Eliaspoor D. Comparison of the effects of low energy laser and Ultrasound in treatment of shoulder myofascial pain syndrome: a randomized single-blinded clinical trial. *Eur J Phys Rehabil Med* 2011;47:381-90

Murtezani A, Hundozi H, Orovcane N, Sllamniku S, Osmani T. A comparison of high intensity exercise and passive modalities for the treatment of workers with chronic low back pain: a randomized, controlled trial. *Eur J Phys Rehabil Med* 2011;47:359-66

Eyigor C, Eyigor S, Kivilcim Korkmaz O. Are intra-articular corticosteroid injections better than conventional TENS in treatment of rotator cuff tendinitis in the short run? A randomized study. *Eur J Phys Rehabil Med* 2011;47:315-24

Hill JC, Whitehurst DGT, Lewis M, Bryan S, Dunn KM, Foster NE, Konstantinou K, Main CJ, Mason E, Somerville S, Sowden G, Vorhora K, Hay EM. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomised controlled trial. *Lancet* 2011;378:1560-71

Hill JC, Dunn KM, Lewis M, Mullis R, Main CJ, Foster NE, Hay EM. A primary care back pain screening tool: indentifying patient subgroups for initial treatment. *Arthritis Rheum US (Arthrit Care Res)* 2008;59:632-41

Wen CP, Wai JPMW, Tsai MK, Yang YC, Cheng TYD, Lee MC, Chan HT, Tsao CK, Wu SPTX. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011;378:1244-53

Marinus J, Moseley GL, Birklein F, Baron R, Maihöfner C, Kingery WS, van Hilten JJ. Clinical features and pathophysiology of complex regional pain syndrome. *Lancet Neurol* 2011;10:637-48

Liebs TR, Herzberg W, Roth-Kroeger M, Rütther W, Hassenpflug J. Women recover faster than men after standard knee arthroplasty. *Clin Orthop Relat Res* 2011;469:2855-65

Kim YH, Choi Y, Kim JS. Comparison of a standard and a gender-specific posterior cruciate-substituting high-flexion knee prosthesis. *J Bone Joint Surg Am* 2010;92:1911-20

Parsley BS, Bertolusso R, Harrington M, Brekke A, Noble PC. Influence of gender on age of treatment with TKA and functional outcome. *Clin Orthop Relat Res* 2010;468(7):1759-64

Kim YH, Choi Y, Kim JS. Comparison of standard and gender-specific posterior-cruciate-retaining high-flexion total knee replacements. *J Bone Joint Surg Br* 2010;92-B:639-45

Bae J, Kong K, Byl N, Tomizuka M. A mobile gait monitoring system for gait analysis. 2009 IEEE, 11th International Conference on Rehabilitation Robotics Kyoto International Conference Center, Japan, June 23-26, 2009. IEEE 2009, 978-1-4244-3789-4/09/\$25.00

Senanayake C, Senanayake SMNA. Human assisted tools for gait analysis and intelligent gait phase detection. 2009, Conference on Innovative Technologies in Intelligent Systems and Industrial Applications (CITSIA 2009) Monash University, Sunway campus, Malaysia, 25th & 26th July 2009. IEEE 2009, 978-1-4244-2887-8/09/\$25.00

Van Dijk GM, Veenhof C, Spreauwenberg P, Coene N, Burger BJ, van Schaardenburg D, van den Ende CH, Lankhorst GJ, Dekker J, on behalf of the CARPA Study Group. Prognosis of limitations in

- activities in osteoarthritis of the hip or knee: a 3-year cohort study. *Arch Phys Med Rehabil* 2010;91:58-66
- Coulter CL, Weber JM, Scarvell JM. Group physiotherapy provides similar outcomes for participants after joint replacement surgery as 1-to-1 physiotherapy: a sequential cohort study. *Arch Phys Med Rehabil* 2009;90:1727-33
- Golomb MR, McDonald BC, Warden SJ, Yonkman J, Saykin AJ, Shirley B, Huber M, Rabin B, AbdelBaky M, Nwosu ME, Barkat-Masih M, Burdea GC. In-home virtual reality videogame telerehabilitation in adolescents with hemiplegic cerebral palsy. *Arch Phys Med Rehabil* 2010;91:1-8
- Kim YH, Choi Y, Kim JS. Comparison of a standard and a gender-specific posterior cruciate-substituting high-flexion knee prosthesis. *J Bone Joint Surg Am* 2010;92:1911-20
- Parsley BS, Bertolusso R, Harrington M, Brekke A, Noble PC. Influence of gender on age treatment with TKA and functional outcome. *Clin Orthop Relat Res* 2010; DOI 10.1007/s11999-010-1348-y
- Dalury DF, Mason JB, Murphy JA, Adams MJ. Analysis of the outcome in male and female patients using a unisex total knee replacement system. *J Bone Joint Surg Br* 2009;91-B:357-60
- Walker FO. Assessing muscle function at the bedside: can we do better? *Muscle Nerve* 2010;42:466-8
- Brooks NE, Cadena SM, Vannier E, Cloutier G, Carambula S, Myburgh K, Roubenoff R, Castandeda-Sceppa C. Effects of resistance exercise combined with essential amino acid supplementation and energy deficit on markers of skeletal muscle atrophy and regeneration during bed rest and active recovery. *Muscle Nerve* 2010; DOI 10.1002/mus.21780
- Padua L, Di Pascale A, Pazzaglia C, Liotta GA, Librante A, Mondelli M. Systematic review of pregnancy-related carpal tunnel syndrome. *Muscle Nerve* 2010;41:697-702
- Duffell LD, Rowleson AM, De N. Donaldson N, Harridge SDR, Newham DIJ. Effects of endurance and strength-directed electrical stimulation training on the performance and histological properties of paralyzed human muscle: a pilot study. *Nerv Muscle* 2010;42:756-63
- McDonald CM, Henricson EK, Han JJ, Abresch RT, Nicorici A, Atkinson L, Elfring GL, Reha A, Miller LL. The 6-minute walk test in Duchenne/Becker muscular dystrophy: longitudinal observations. *Muscle Nerv* 2010; DOI 10.1002/mus.21808
- Kallenberg LAC, Hermens HJ. Motor unit properties of biceps brachii during dynamic contractions in chronic stroke patients. *Muscle Nerve* 2010; DOI 10.1002/mus.21803
- Lin H, Hou C, Chen D. Modified C7 neurotization for the treatment of obstetrical brachial plexus palsy. *Muscle Nerve* 2010;42:764-8
- McNeely ML, Campbell KL, Courneya KS, Mackey JR. Effect of acute exercise on upper-limb volume in breast cancer survivors: a pilot study. *Physiother Can* 2009;61:244-51
- Kilbreath SL, Ward LC, Lane K, McNeely M, Dylke ES, Refshauge KM, McKenzie D, Lee MJ, Peddle C, Battersby KJ. Effect of air travel on lymphedema risk in women with history of breast cancer. *2010;120:649-54*
- Proding B, Weise AP, Shaw L, Stamm TA. A Delphi study on environmental factors that impact work and social life participation of individuals with multiple sclerosis in Austria and Switzerland. *Disabil Rehabil* 2010;32:183-95

Badger CMA, Peacock JL, Mortimer PS. A randomized, controlled, parallel-group clinical trial comparing multilayer bandaging followed by hosiery versus hosiery alone in the treatment of patients with lymphedema of the limb. *Cancer* 2000;88:2832-7

Vignes S, Porcher R, Arrault M, Dupuy A. Long-term management of breast cancer-related lymphedema after intensive decongestive physiotherapy. *Breast Cancer Res Treat* 2007;101:285-90

Shaw C, Mortimer P, Judd PA. A randomized controlled trial of weight reduction as a treatment for breast cancer-related lymphedema. *Cancer* 2007;110:1868-74

Berschin G, Sommer HM. Zur Bedeutung der Körperhaltung auf Energietransmission und -absorption beim Vibrationstraining. *Sportverl Sportschad* 2010;24:36-39

Naal FD, Impellizzeri FM, Wasmer M, Mannion AF, Leunig M. Patientenorientierter Schulthess-Hüftscore (5 Items) zur Evaluation von Koxarthrose- und Hüftarthroplastikpatienten. *Orthopäde* 2010; DOI 10.1007/s00132-010-1611-7

Naal FD, Sieverding M, Impellizzeri FM, von Knoch F, Mannion AF, Leunig M. Reliability and validity of the Cross-Culturally Adapted German Osford Hip Score. *Clin Orthop Relat Res* 2009;467:952-7

Ruedl G, Sommersacher R, Woldrich T, Pocecco E, Hotter B, Nachbauer W, Burtscher M. Wer trägt einen Skihelm? Helmtragequote auf österreichischen Skipisten in Abhängigkeit verschiedener Einflussfaktoren. *Sportverl Sportschad* 2010;24:27-30

May AM, van Weert E, Korstjens I, Hoekstra-Weebers JE, van der Schans CP, Zonderland ML, Mesters I, van den Borne B, Ros WJ. Monitoring training progress during exercise training in cancer survivors: a submaximal exercise test as an alternative for a maximal exercise test? *Arch Phys Med Rehabil* 2010;91:351-7

Rubi M, Renom F, Ramis F, Medinas M, Centeno MJ, Górriz M, Crespi E, Marin B, Soriano JB. Effectiveness of pulmonary rehabilitation in reducing health resources use in chronic obstructive pulmonary disease. *Arch Phys Med Rehabil* 2010;91:364-8

Rollman A, Naeije M, Visscher CM. The reproducibility and responsiveness of a patient-specific approach: a new instrument in evaluation of treatment of temporomandibular disorders. *J Orofac Pain* 2010;24:101-5

Cakit BD, Nacir B, Genc H, Saracoglu M, Karagöz A, Erdem HR, Ergün U. Cycling progressive resistance training for people with multiple sclerosis: a randomized controlled study. *Am J Phys Med Rehabil* 2010; DOI: 10.1097/PHM.0b013e3181d3e71f

Hébert-Blouin MN, Amrami KK, Wang H, Skinner JA, Spinner RJ. Tibialis anterior branch involvement in fibular intraneural ganglia. *Muscle Nerve* 2010;41:524-32

Cammarata ML, Dhaher YY. Evidence of gender-specific motor templates to resist valgus loading at the knee. *Muscle Nerve* 2010;41:614-23

Serafin A, Molin J, Márquez M, Blasco E, Vidal E, Foradada L, Anor S, Rabanal RM, Fondevila D, Bosch F, Pumarola M. Diabetic neuropathy: electrophysiological and morphological study of peripheral nerve degeneration and regeneration in transgenic mice that express IFN β in β Cells. *Muscle Nerve* 2010;41:630-41

Bernsen RAJAM, de Jager AEJ, Kuijter W, van der Meché FGA, Suurmeijer TPBM. Psychosocial dysfunction in the first year after Guillain-Barré Syndrome. *Muscle Nerve* 2010; 41:533-9

McDonald CM, Henricson EK, Han JJ, Abresch RT, Nicorici A, Elfring GL, Atkinson L, Reha A, Hirawat S, Miller LL. The 6-minute walk test as a new outcome measure in duchenne muscular dystrophy. *Muscle Nerve* 2010;41:500-10

- Bayrak AO, Bayrak IK, Turker H, Elmali M, Nural MS. Ultrasonography in patients with ulnar neuropathy at the elbow: comparison of cross-sectional area and swelling ratio with electrophysiological severity. *Muscle Nerve* 2010;41:661-6
- Malladi N, Micklesen PJ, Hou J, Robinson LR. Correlation between the combined sensory index and clinical outcome after carpal tunnel decompression: a retrospective review. *Muscle Nerve* 2010;41:453-7
- Gigo-Benato D, Russo TL, Geuna S, Dominigues NRSR, Salvini TF, Parizotto NA. Electrical stimulation impairs early functional recovery and accentuates skeletal muscle atrophy after sciatic nerve crush injury in rats. *Muscle Nerve* 2010;41:685-93
- Gschiel B, Kager H, Pipam W, Weichart K, Likar R. Analgetische Effizienz von transkutaner elektrischer Nervenstimulation (TENS-Therapie) bei Patienten mit GORarthrose. *Schmerz* 2010; DOI 10.1007/s00482-010-0957-4
- Burke TN, Franca FJR, de Meneses SRF, Cardoso VI, Marques AP. Postural control in elderly persons with osteoporosis: Efficacy of an intervention program to improve balance and muscle strength: a randomized controlled trial. *Am J Phys Med Rehabil* 2010;89:549-56
- Meier S, Neubauer E, Schiltenswolf M. Messung von Behandlungserfolg bei chronischen Rückenschmerzen. *Schmerz* 2009; DOI 10.1007/s00482-008-0731-z
- Frobell RB, Roos EM, Roos HP, Ranstam J, Lohmander S. A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med* 2010;363:331-42
- Edwards BJ, Song J, Dunlop DD, Fink HA, Cauley JA. Functional decline after incident wrist fractures - Study of Osteoporotic Fractures: prospective cohort study. *BMJ* 2010;341:c3324
- Wang C, Schmid CH, Rones R, Kalish R, Yin J, Goldenberg DL, Lee Y, McAlindon T. A randomized trial of Tai Chi for fibromyalgia. *N Engl J Med* 2010;363:743-54
- Spranger M, Deppe W. Assessments in der neurologischen Rehabilitation von Kindern und Jugendlichen. *Neurol Rehabil* 2007;13:15-23
- Depedibi R, Ünlü E, Cevikol A, Akkaya T, Cakc A, Cerekci R, Köse G, Ünlüsoy D. Ultrasound-guided botulinum toxin type A injection to the iliopsoas muscle in the management of children with cerebral palsy. *Neuro Rehabil* 2008;23:199-205
- Stearns GE, Burtner P, Keenan KM, Qualls C, Phillips J. Effects of constraint-induced movement therapy on hand skills and muscle recruitment of children with spastic hemiplegic cerebral palsy. *Neuro Rehabil* 2009;24:95-108
- Andersen LB, Lawlor DA, Cooper AR, Froberg K, Anderssen SA. Physical fitness in relation to transport to school in adolescents: the Danish youth and sports study. *Scand J Med Sci Sports* 2009;19:406-11
- Bartl R, Gradinger R. Aktuelle Diagnostik und Therapie der Osteoporose auf der Basis der „European Guidance 2008“. *Orthopädie* 2009;38:365-80
- Michalski D, Zweynert U, Kittel J, Hinz A. Schmerzempfinden und -verhalten während der orthopädischen Rehabilitation. *Schmerz* 2009; DOI 10.1007/s00482-009-0778-5
- Rollnik JD. Der Barthel-Index als Verweildauer-Prädiktor in der neurologischen Rehabilitation. *Rehabilitation* 2009;48:91-4

Greiner S, Herrmann S, Gerhardt C, Scheibel M. Klassifikation und Diagnostik der instabilen Schulter. Orthopädie 2009;38:6-15

Gruther W, Benesch T, Zorn C, Paternostro-Slugla T, Quittan M, Fialka-Moser V, Spiss C, Kainberger F, Crevenna R. Muscle wasting in intensive care patients: ultrasound observation of the M. quadriceps femoris muscle layer. J Rehabil Med 2007; DOI: 10.2340/16501977-0139

Sanatani M, Schreier G, Stitt L. Level and direction of hope in cancer patients: an exploratory longitudinal study. Support Care Cancer 2007; DOI: 10.1007/s00520-007-0336-6

Zorn C, Szekeres T, Keilani M, Fialka-Moser V, Crevenna R. Effects of neuromuscular electrical stimulation of the knee extensor muscles on muscle soreness and different serum parameters in young male athletes: preliminary data. Br J Sports Med 2007;41:914-16

Keilani MY, Posch M, Zorn C, Knötig M, Pircher M, Quittan M, Fialka-Moser V, Jirasek U, Crevenna R. Vergleichende Analyse der Messwtergebnisse von zwei Dynamometern zur Messung der isokinetischen Kraft der kniegelenksbewegenden Muskulatur. Phys Med Rehab Kuror 2007;17:327-33

Hooten WM, Townsend CO, Decker PA. Gender differences among patients with fibromyalgia undergoing multidisciplinary pain rehabilitation. Pain Medicine 2007;8:624-32

Kool JP, Oesch PR, De Bie RA. Predictive tests for non-return to work in patients with chronic low back pain. Eur Spine J 2002;11:258-66

Claus A, Hides J, Moseley GL, Hodges P. Sitting versus standing: does the intradiscal pressure cause disc degeneration of low back pain? J Electromyogr Kines 2008;18:550-8

Bachmann S, Eisenring M, Kool J, Oesch P, Michel BA, Knüsel O. Rehabilitations- und Folgekosten bei Patienten mit chronischen Rückenschmerzen. Phys Med Rehab Kuror 2008;18:181-8

Adams J, Burrige J, Mullee M, Hammond A, Cooper C. The clinical effectiveness of static resting splints in early rheumatoid arthritis: a randomized controlled trial. Rheumatol 2008;47:1548-53

Peters-Veluthamaningal C, Winters JC, Groenier KH, Meyboom-de Jong B. Corticosteroid injections effective for trigger finger in adults in general practice: a double-blinded randomised placebo controlled trial. Ann Rheum Dis 2008;67:1262-6

Güler-Yüksel M, Bijsterbosch J, Goekoop-Ruiterman YPM, de Vries-Bouwstra JK, Hulsmans HMJ, de Beus WM, KH Han, Breedveld FC, Dijkmans BAC, Allaart CF, Lems WF. Changes in bone mineral density in patients with recent onset, active rheumatoid arthritis. Ann Rheum Dis 2008;67:823-8

Schädler, Kool, Lüthi, Marks, Pfeffer, Oesch, Wirz. Assessments in der Neurorehabilitation. Functional Ambulation Categories (FAC). Verlag Hans Huber 2006.

Scherfer E, Bohls C, Freiburger E, Heise KF, Hogan D. Manual Berg Balance Scale (BBS). Schädler, Kool, Lüthi, Marks, Pfeffer, Oesch, Wirz. Assessments in der Neurorehabilitation. Verlag Hans Huber 2006.

Bensoussan L, Viton JM, Barotsis N, Delarque A. Evaluation of patients with gait abnormalities in physical and rehabilitation medicine settings. J Rehabil Med 2008;40:497-507

Thomas NE, Williams DRR. Inflammatory factors, physical activity, and physical fitness in young people. Scand J Med Sci Sports 2008;18:543-56

Sapsford RR, Richardson CA, Maher CF, Hodges PW. Pelvic floor muscle activity in different sitting postures in continent and incontinent women. Arch Phys Med Rehabil 2008;89:1741-7

Davis R, Sparrow O, Cosendai G, Burridge JH, Wulff C, Turk R, Schulman. Poststroke upper-limb rehabilitation using 5 to 7 inserted microstimulators: implant procedure, safety, and efficacy for restoration of function. *Arch Phys Med Rehabil* 2008;89:1907-12

Schultheis A, Reichwein F, Nebelung W. Die eingesteifte Schulter. *Orthopäde* 2008;11:2065-72

Peul WC, van den Hout WB, Brand R, Thomeer TWM, Koes BW. Prolonged conservative care versus early surgery in patients with sciatica caused by lumbar disc herniation: two year results of a randomised controlled trial. *BMJ* 2008;336:1355-8

Kirkley A, Birmingham TB, Litchfield RB, Giffin JR, Willits KR, Wong CJ, Feagan BG, Donner A, Griffin SH, D'Ascanio LM, Pope JE, Fowler PJ. A Randomized trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med* 2008;359:1097-107

Englund M, Guermazi A, Gale D, Hunter DJ, Aliabadi P, Clancy M, Felson T. Incidental meniscal findings on knee MRI in middle-aged and elderly persons. *N Engl J Med* 2008;359:1108-15

Van den Hout WB, Peul WC, Koes BW, Brand R, Kievit J, Thomeer TWM. Prolonged conservative care versus early surgery in patients with sciatica from lumbar disc herniation: cost utility analysis alongside a randomised controlled trial. *BMJ* 2008;336:1351-4

Maki BE, Cheng KCC, Mansfield A, Scovil CY, Perry SD, Peters AL, McKay S, Lee T, Marquis A, Corbeil P, Fernie GR, Liu B, McIlroy WE. Preventing falls in older adults: new interventions to promote more effective change-in-support balance reactions. *J Electromyogr Kines* 2008;18:243-54

Maurer C, Mergner T, Lücking CH, Becker W. Adaptive changes of saccadic eye-head coordination resulting from altered head posture in torticollis spasmodicus. *Brain* 2001;124:413-26

Candia V, Schäfer M, Taub E, Rau H, Altenmüller E, Rockstroh B, Elbert T. Sensory motor retuning: a behavioral treatment for focal hand dystonia of pianists and guitarists. *Arch Phys Med Rehabil* 2002;83:1342-8

Altenmüller E. Focal dystonia: advances in brain imaging and understanding of fine motor control in musicians. *Hand Clin* 2003;19:1-16

Prof. Dr. Becker. Okulomotorik und vestibuläres System. Klinikbericht der Abteilung für Neurologie der Universität Ulm. <http://www.uni-ulm.de/klinik/expneuro/klinikbericht-9699-07.html>

Arnson Y, Amital H, Shoenfeld Y. Vitamin D and autoimmunity: new aetiological and therapeutic considerations. *Ann Rheum Dis* 2007;66:1137-42

Roddy E, Zhang W, Doherty M. Is gout associated with reduced quality of life? A case-control study. *Rheumatol* 2007;46:1441-4

Valdes AM, Doherty M, Spector TD. The additive effect of individual genes in predicting risk of knee osteoarthritis. *Ann Rheum Dis* 2007;67:124-7

Lubrano E, Angelo SD, Parsons WJ, Corbi G, Ferrara N, Rengo F, Olivieri I. Effectiveness of rehabilitation in active ankylosing spondylitis assessed by the ASAS response criteria. *Rheumatol* 2007;46:1672-5

Vasodilator iontophoresis – a possible new therapy for digital ischaemia in systemic sclerosis? *Rheumatol* 2008;47:76-9

Huber W, Hofstaetter JG, Hanslik-Schnabel B, Posch M, Wurnig C. The German version of the Oxford shoulder score – cross-cultural adaptation and validation. *Arch Orthop Trauma Surg* 2004;124:531-6

Dawson J, Rogers K, Fitzpatrick R, Carr A. The Oxford shoulder score revisited. Arch Orthop Trauma Surg. DOI 10.1007/s00402-007-0549-7

Elsner G. Anerkennung von Berufskrankheiten im europäischen Vergleich. Bundesgesundheitsbl Gesundheitsforsch Gesundheitsschutz 2008;51:281-6

Coster W, Alkunkal Khetani M. Measuring participation of children with disabilities: Issues and challenges. Disabil Rehabil 2008;30:369-48

Yavuzer G, Selles R, Sezer N, Sütbeyaz S, Bussmann JB, Köseoglu F, Atay MB, Stam HJ. Mirror therapy improves hand function in subacute stroke: a randomized controlled trial. Arch Phys Med Rehabil 2008;89:393-8

Meserve BB, Cleland JA, Boucher TR. A meta-analysis examining clinical test utilities for assessing meniscal injury. Clin Rehabil 2008;22:143-61

Myint JM, Yuen GF, Yu TK, Kng CP, Wong AM, Chow KK, Li HC, Wong CP. A study of constraint-induced movement therapy in subacute stroke patients in Hong Kong. Clin Rehabil 2008 22:112-24

Costa BR, Vieira ER. Stretching to reduce work-related musculoskeletal disorders: a systematic review. J Rehabil Med 2008;40:321-8

Zhang JG, Ishikawa-Takata K, Yamazaki H, Morita T, Ohta T. Postural stability and physical performance in social dancers. Gait Posture 2008;27:697-701

Burton MJ, Monsell EM, Rosenfeld RM. Extracts from the Cochrane Library: Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. Otolaryng Head Neck 2008;138:415-7

Jaber JJ, Leonetti JP, Lawrason AE, Feustel PJ. Cervical spine causes for referred otalgia. Otolaryng Head Neck 2008;138:479-85

Raschner C, Lember S, Platzer HP, Patterson C, Hilden T, Lutz M. S3-Check-Evaluierung und Normwertenerhebung eines Tests zur Erfassung der Gleichgewichtsfähigkeit und Körperstabilität. Sportverl Sportschad 2008;22:100-5

Dehner C, Heym B, Maier D, Sander S, Arand M, Elbel M, Hartwig E, Kramer M. Postural control deficit in acute QTF grade II whiplash injuries. Gait Posture 2008;28:113-9

Rose KJ, Burns J, Ryan MM, Ouvrier RA, North KN. Reliability of quantifying foot and ankle muscle strength in very young children. Muscle Nerve 2008;37:626-31

Foley J, Little JW, Vaezy S. Effects of high-intensity focused ultrasound on nerve conduction. Muscle Nerve 2008;37:241-50

Gerner P, Strichartz GR. Sensory and motor complications of local anesthetics. Muscle Nerve 37:421-5

Ginanneschi F, Milani P, Mondelli M, Dominici F, Biasella A, Rossi A. Ulnar sensory nerve impairment at the wrist in carpal tunnel syndrome. Muscle Nerve 2008;183-9

Mirbagheri MM, Tsao CC, Rymer WZ. Changes of elbow kinematics and kinetics during 1 year after stroke. Muscle Nerve 2008;37:387-95

Jackson CE, Barohn RJ, Gronseth G, Pandya S, Herbelin L, the muscle study group. Inclusion body myositis functional rating scale: a reliable and valid measure of disease severity. Muscle Nerve 2008;37:473-6

Roquelaure Y, Ha C, Pelier-Cady MC, Nicolas G, Descatha A, Leclerc A, Raimbeau G, Goldberg M, Imbernon E. Work increases the incidence of carpal tunnel syndrome in the general population. *Muscle Nerve* 2008;37:477-82

Rajaratnam BS, Venketasubramanian N, Kumar PV, Goh JC, Chan YH. Predictability of simple clinical tests to identify shoulder pain after stroke. *Arch Phys Med Rehabil* 2007;88:1016-21

Kang BS, Bang MS, Jung SH. Effects of botulinum toxin A therapy with electrical stimulation on spastic calf muscles in children with cerebral palsy. *Am J Phys Med Rehabil* 2007;86:000-000

Floman Y, Smorgick Y, Rand N, Bar-Ziv J. Inferior vena cava thrombosis presenting as lumbar radiculopathy. *Am J Phys Med Rehabil* 2007;86:000-000

Petersen W, Welp R, Rosenbaum D. Chronic achilles tendopathy. A prospective randomized study comparing the therapeutic effect of eccentric training, the Airheel brace, and a combination of both. *Am J Sports Med* 2007;35: 1659-67

Taki M, Iwata O, Shiono M, Kimura M, Takagishi K. Extracorporeal shock wave therapy for resistant stress fracture in athletes: a report of 5 cases. *Am J Sports Med* 2007;35:1659

Sachs RA, Lin D, Stone ML, Paxton E, Kuney M. Can the need for future surgery for acute traumatic anterior shoulder dislocation be predicted? *Am J Bone Joint Surg* 2007;89:1665-74

Kool J, Bachmann S, Oesch P, Knuesel O, Amberg T, de Bie R, van den Brandt P. Function-centered rehabilitation increases work days in patients with nonacute nonspecific low back pain: 1-year results from a randomized controlled trial. *Arch Phys Med Rehabil* 2007;88:1089-94

Jonsdottir J, Cattaneo D. Reliability and Validity of the dynamic gait index in persons with chronic stroke. *Arch Phys Med Rehabil* 2007;88:1410-5

Vila B, Servera E, Marin J, Diaz J, Giménez M, Komaroff E, Bach J. Noninvasive ventilatory assistance during exercise for patients with kyphoscoliosis: a pilot study. *Am J Phys Med Rehabil* 2007;86:672-7

Chow IHW, Cheing GLY. Comparison of different energy densities of extracorporeal shock wave therapy (ESWT) for the management of chronic heel pain. *Clin Rehabil* 2007;21:131-41

Schädler S. Balance beim Gehen beurteilen. *Physiotherapie* 2006;10:40-1

Cattaneo D, Jonsdottir J, Zocch M, Regola A. Effects of balance exercise on people with multiple sclerosis: a pilot study. *Clin Rehabil* 2007;21:771-81

McCarthy CJ, Callaghan MJ, Oldham JA. Pulsed electromagnetic energy treatment offers no clinical benefit in reducing the pain of knee osteoarthritis: a systematic review. *BMC Musculoskeletal Disorders* 2006;7:51

Vassiliou T, Kaluza G, Putzke C, Wulf H, Schnabel M. Physical therapy and active exercises – an adequate treatment for prevention of late whiplash syndrome? Randomized controlled trial in 200 patients. *Pain* 2006;

Murphy SB, Tannast M. Herkömmliche vs. Minimal-invasive Hüftendoprothetik. Eine prospektive Studie über Rehabilitation und Komplikationen. *Orthopäde* 2006;10.2007/s00132-006-0969-z

Schilling AM, Heidenreich JO, Wolf KJ. Fallgruben und Besonderheiten der modernen bildgebenden Diagnostik der oberen Halswirbelsäule. *Orthopäde* 2006;35:237-43

Treaster D, Marras WS, Burr D, Sheedy JE, Hart D. Myofascial trigger point development from visual and postural stressors during computer work. *J Electromyogr Kinesiol* 2006;16:115-24

Chiodo A, Goodmurphy C, Haig A. Cadaveric study of methods for subscapularis muscle needle insertion. *Am J Phys Med Rehabil* 2005;84:662-5

Yaar I. The logical choice of muscles for the needle-EMG evaluation of lumbosacral radiculopathy. *J Electromyogr Kinesiol* 2006;16:205-13

Hambly K, Bobic V, Wondrasch B, Van Assche D, Marlovits S. Autologous chondrocyte implantation postoperative care and rehabilitation. *Am J Sports Med* 2006;34:1020-38

Turkel CC, Bowen B, Liu J, Brin MF. Pooled analysis of the safety of Botulinum Toxin Type A in the treatment of poststroke spasticity. *Arch Phys Med Rehabil* 2006;87:786-92

Badke MB, Boissonnault WG. Changes in disability following physical therapy intervention for patients with low back pain: dependence on symptom duration. *Arch Phys Med Rehabil* 2006;87:749-56

Medlicott MS, Harris SR. A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder. *Phys Ther* 2006;86:955-73

Thuret S, Moon LDF, Gage FH. Therapeutic interventions after spinal cord injury. *Nature* 2006;7:62843

Fejer R, Hartvigsen J, Kyvik KO. Heritability of neck pain: a population-based study of 33794 Danish twins. *Rheumatology* 2006;45:589-94

Pollard LC, Choy EH, Gonzalez J, Khoshaba B, Scott DL. Fatigue in rheumatoid arthritis reflects pain, not disease activity. *Rheumatology* 2006;45:885-9

Van de Port IGL, Kwakkel G, Schepers VPM, Lindeman E. Predicting mobility outcome on year after stroke: a prospective cohort study. *J Rehabil Med* 2006;38:218-23

Graham N, Gross AR, Goldsmith C, Cervical Overview Group. Mechanical traction for mechanical neck disorders: a systematic review. *J Rehabil Med* 2006;38:145-52

Blumenthal S, Herskovitz S, Verghese J. Carpal tunnel syndrome in older adults. *Muscle Nerve* 2006;34:78-83

Rankin G, Stokes M, Newham DJ. Abdominal muscle size and symmetry in normal subjects. *Muscle Nerve* 2006;34:320-6